

Impact of UV Light Disinfection on CLABSI and CAUTI Rates in a Major Medical Teaching Hospital's MICU

Susan Hanrahan, MS, CIC, CRCST, Infection Prevention Manager JSUMC, Chad Haraschak, BS, Environmental Services, Crothall Healthcare, Gary Buckalew, RN, MSN, CCRN, Nurse Manager MICU/CCU JSUMC

Abstract

Ultra Violet-C (UV-C) light disinfection was consistently applied to the MICU/CCU from 03/2015 through 07/2015. UV-C light disinfection of the unit was an adjunct to the routine chemical disinfection program already in existence. NHSN SIR metrics were used to measure the impact of UV-C light disinfection. The MICU/CCU experienced a statistically significant reduction in CAUTI incidence and a measurable reduction in CLABSI incidence, although not statistically significant for the 5 month study period that UV-C light disinfection was consistently used. When UV-C light disinfection was withdrawn and delivered as a less vigorous intervention, CAUTI and CLABSI increased, although not statistically significant.

Introduction

Previous authors have established that UV light disinfection has been observed to decrease health care acquired infections (HAIs) in both acute care and long term care facilities^{1,2,3}. Specifically, UV light disinfection utilizing UV-C radiation was effective at decreasing the bioburden of MRSA, C. diff, Enterococcus and Acinetobacter in these studies. UV light systems that can deliver UV-C radiation at approximately 254 nm have been documented to eradicate pathogens by denaturing their DNA. UV light systems emitting at this wavelength are capable of breaking down the DNA nucleotide bonds, thus rendering microorganisms incapable of replicating.

Objective

Here we evaluated the efficacy of an automated UV-C triple emitter system, which has been designed to accomplish room disinfection more effectively and in less time. CAUTI and CLABSI rates were evaluated in the MICU/CCU before and after the consistent application of UV-C light disinfection as an Infection Control and Housekeeping intervention. Initially brought on board to JSUMC as a test device through the Crothall Healthcare Products Committee, the UV-C triple emitter matured from a "test and trial" program into a "daily addition" disinfection process in the MICU/CCU. UV-C light disinfection was used in conjunction with the chemical disinfection program deployed by Housekeeping.

Methods

Each discharge room was effectively chemically disinfected, curtain changed, terminally cleaned, and treated with UV-C light. UV-C light was available for use in the MICU/CCU as needed for every discharge during the study period. Protocols for MICU/CCU, Housekeeping, and Bed Management impacting admission and discharge were altered to accommodate this technology. Working collaboratively with MICU/CCU staff regarding these processes, proved to be instrumental in outcomes. The UV-C light disinfection system was consistently applied to the MICU/CCU from 03/2015 through 07/2015 in addition to the housekeeping program already in place. NHSN SIR metrics were used to measure the impact of UV-C light disinfection on CAUTI and CLABSI rates before and after the intervention.

Results

Prior to the study period, the 2014 YTD CAUTI data was unfavorably statistically significant (SIR=1.93, $p=0.00$, SIR95CI=1.214, 2.933) (Figure 1). During the study period when UV-C light disinfection was consistently used, the MICU experienced a statistically significant reduction in CAUTI incidence (SIR =0.00, $p=0.01$, SIR95CI =, 0.673) and a measurable reduction in CLABSI incidence (SIR=0.036, $p=0.29$, SIR95CI = 0.018, 1.763). After the study period when UV-C light disinfection was less consistently applied, CAUTI and CLABSI increased, although not statistically significant (CAUTI SIR=1.44, $p=0.41$, SIR95CI=0.527, 3.190; CLABSI SIR=1.81, $p=0.21$, SIR95CI=0.662, 4.004).

Conclusion

Prior to the new UV-C light disinfection program at Jersey Shore University Medical Center (JSUMC), the Infection Control Committee was striving to reduce the incidence of both CAUTI and CLABSI for over a year. A CAUTI Action Plan had been in place since fourth quarter 2014, secondary to unfavorable year end data (Figure 1). Several nursing improvements to patient care had already been implemented prior to the addition of the triple emitter UV-C system. The addition of the automated UV-C disinfection system further reduced both CAUTI and CLABSI incidence when applied consistently in the MICU.

Environmental contamination of the patient care area is a potential contributing factor to HAI incidence. Several studies demonstrate how the environment of care is integral to preventing transmission of pathogenic organisms¹. Chemical disinfection has been the gold standard for healthcare, heavily relying on quaternary ammonium and bleach based disinfectant preparations. Previous studies have demonstrated that less than 50% of contaminated surfaces actually make contact with the chemical disinfectant¹. Establishing a housekeeping program with reliable chemical disinfection efficacy can be challenging. It is speculated that the dedicated UV-C light disinfection program tested in the MICU/CCU was decreasing bioburden for safer delivery of patient care related to CAUTI and CLABSI prevention, more so than just chemical disinfection alone.

References

- Vianna P, Dale C, Simmons S, Stibich M, Licitra C. Impact of pulsed xenon ultraviolet light on hospital-acquired infection rates in a community hospital. Am J Infect Control 2016;44:299-303.
- Miller R, Simmons S, Dale C, Stachowiak J, Stibich M. Utilization and impact of a pulsed-xenon ultraviolet room disinfection system and multidisciplinary care team on *Clostridium difficile* in a long-term acute care facility. Am J Infect Control 2016;43:1350-5.
- Napolitano N, Mahapatra T, Weiming T. The effectiveness of UV-C radiation for facility-wide environmental disinfection to reduce health care-acquired infections. Am J Infect Control 2016;43:1342-6.

Figure 1. 2015 MICU Infection Control Dashboard

		Actual Rate												Actual Rate			Actual Rate			2015 YTD SIR					
2015 Mandated Reported Measures		2014												3rd Q			4th Q			2015 YTD SIR					
Goal SIR <=1		YTD 2014		1st Q			2nd Q			3rd Q			4th Q			2015 YTD SIR									
		Rate	SIR	Jan	Feb	Mar	Rate	1st Q SIR	Apr	May	Jun	Rate	SIR	Jul	Aug	Sep	Rate	3rd Q SIR	Oct	Nov	Dec	Rate	SIR	2015 YTD SIR	
18	MICU	1.6	0.81	2.9	0.0	2.8	0.0	1.00	0.0	0.0	0.0	0.0	0.0	0.0	7.4	3.3	3.8	1.88	7.0	0.0	0.0	2.5	1.02	0.98	
19	CLABSI	5	5	1	0	1	2	2	0	0	0	0	0	0	2	1	3	3	2	0	0	2	2	2	7
20		3086	6	349	0.99	357	1005	2	283	286	250	819	2	223	389	305	797	2	286	245	278	810	2	7	
21			$p=0.08$					$p=0.00$					$p=0.19$					$p=0.29$					$p=0.70$	$p=0.91$	
22	CAUTI	3.9	1.89	0.0	4.6	0.0	0.0	0.67	0.0	0.0	0.0	0.0	0.0	0.0	2.7	3.0	1.9	0.94	5.9	0.0	2.6	2.9	1.44	0.72	
23		20	20	0	2	2	2	0	0	0	0	0	0	0	1	1	2	2	2	0	2	3	3	2	
24		5172	41	470	458	581	1489	3	468	444	361	1273	3	371	364	330	1065	2	337	328	378	1043	2	10	
25			$p=0.00$					$p=0.63$					$p=0.08$					$p=0.00$					$p=0.11$	$p=0.00$	

Decreased incidence

Increased incidence

Key Terms:

CAUTI = Catheter Associated Urinary Tract Infection
 CLABSI = Central Line Associated Blood Stream Infection
 NHSN = National Healthcare Safety Network
 SIR = Standardized Infection Ratio

Acknowledgements

Adam Buchaklian PhD